



# City Research Online

## City, University of London Institutional Repository

---

**Citation:** Elliott, D., McKinley, S., Alison, J., Aitken, L. M. ORCID: 0000-0001-5722-9090 and King, M. (2006). Study protocol: Home-based physical rehabilitation for survivors of a critical illness [ACTRN12605000166673]. *Critical Care*, 10(3), R90. doi: 10.1186/cc4949

This is the published version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/25921/>

**Link to published version:** <http://dx.doi.org/10.1186/cc4949>

**Copyright and reuse:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

## Research

## Open Access

**Study protocol: Home-based physical rehabilitation for survivors of a critical illness [ACTRN12605000166673]**Doug Elliott<sup>1</sup>, Sharon McKinley<sup>2</sup>, Jennifer A Alison<sup>3</sup>, Leanne M Aitken<sup>4</sup> and Madeleine T King<sup>5</sup><sup>1</sup>Faculty of Nursing, Midwifery and Health, University of Technology, Sydney, New South Wales, Australia<sup>2</sup>Critical Care Professorial Unit, University of Technology, Sydney & Northern Sydney Central Coast Area Health Service, New South Wales, Australia<sup>3</sup>School of Physiotherapy, Faculty of Health Sciences, The University of Sydney, New South Wales, Australia<sup>4</sup>Princess Alexandra Hospital & Griffith University, Queensland, Australia<sup>5</sup>Centre for Health Economics, Research & Evaluation, University of Technology, Sydney, New South Wales, AustraliaCorresponding author: Doug Elliott, [doug.elliott@uts.edu.au](mailto:doug.elliott@uts.edu.au)

Received: 9 May 2006 Accepted: 15 May 2006 Published: 22 Jun 2006

*Critical Care* 2006, **10**:R90 (doi:10.1186/cc4949)This article is online at: <http://ccforum.com/content/10/3/R90>© 2006 Elliott *et al.*; licensee BioMed Central Ltd.This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**Abstract**

**Introduction** Numerous primary studies and several review papers have highlighted delayed physical and psychological recovery for survivors of critical illness, often beyond 6 months after discharge. This randomized controlled trial with blinded assessment aims to test the effects of an 8-week, home-based, individually tailored physical rehabilitation programme on physical and psychological recovery for survivors of a critical illness after discharge from hospital.

**Method** Participants are survivors of a critical illness discharged from nine intensive care units (ICUs) in Australia, who are aged 18 years or older, in an ICU longer than 48 hours, discharged home to self-care or carer (non-institutional care), able to participate in physical rehabilitation, and within the hospitals' local geographical areas for home visits. The study is based in participants' home environments. Blinded assessments at weeks 1, 8 and 26 after hospital discharge examine physical functioning, exercise capacity, health-related quality of life and psychological well being. The intervention is graded, individualized endurance and strength training prescribed by a pulmonary rehabilitation physiotherapist over an 8-week period, with three home visits, five follow-up phone calls, and a printed

exercise manual supporting the training. Initial focus is on lower limb exercises and walking, with warm-up stretches, and progresses to the addition of core stabilization and upper limb exercises.

**Results** The burden of a critical illness is well documented. This novel study will determine whether a home-based physical rehabilitation programme improves the recovery trajectory for survivors of critical illness. The projected sample size of 200 patients aims to detect a clinically important 10% improvement in physical functioning. The study will also examine whether other important physical and psychological measures are improved.

**Conclusion** This multicentre, randomized controlled trial will examine outcomes that are meaningful to patients, their family and society, namely functional ability and well being. The study will also target a health problem that is likely to increase as the population ages. If the programme is effective, it will provide a model that can be easily adapted and adopted by existing primary care or community services to improve the recovery of individuals following critical illness.

**Introduction**

Examination of patient outcomes following a critical illness, over and above survival, has become an important topic of critical care research, particularly over the last decade [1,2]. Although continuing patient follow up by the intensive care unit (ICU) team following ICU discharge is not a traditional routine practice, this lack of continuity is now being questioned [3].

The development of critical care outreach or liaison teams is evident, particularly in the UK [4] and to a lesser extent in Australia [5], with a focus on early detection of clinical deterioration. Importantly, however, systematic follow up with physical rehabilitation interventions for survivors of critical illness with physical debilitation is not routinely provided, particularly after hospital discharge, although there are exceptions [6].

DASS = Depression, Anxiety, and Stress Scale; HRQOL = health-related quality of life; ICU = intensive care unit; IES = Impact of Events Scale; 6 MWT = 6 minute walk test; PF = Physical Functioning (subscale); RCT = randomized controlled trial; 8 RM = eight-repetition maximum; SF-36 = Short-Form-36 version 2.

A critical illness requiring admission to a general ICU affects over 130,000 adult Australians per year [7]. Although survival rates approximate 86% [7], functional recovery for individuals is often delayed beyond 6 months after hospital discharge [8-10]. There is commonly physical de-conditioning [11] as well as psychological sequelae [12,13], adding to the burden of illness for our society [3]. A number of review papers identified a plethora of observational studies that confirm this delayed recovery [8-10]. The methods and measuring instruments used to explore patient outcomes have evolved from crude mortality and morbidity indicators to more patient-centred concepts such as functional status and health-related quality of life (HRQOL) [9]. Some methodological limitations with these more recent studies remain, including the lack of randomized trials, use of unvalidated measuring instruments and lack of control of extraneous variables [14]. Despite these limitations, it is clear that significant sequelae exist for a substantial proportion of critical illness survivors. However, there is a lack of evidence for the benefit of any specific service provision for recovering survivors from critical illness [3,15], with only a few published interventional studies available [6,16]. None have tested the effect of home-based rehabilitation on patient recovery.

We propose that a focused home-based approach to physical rehabilitation, in addition to usual community-based health services, will improve the HRQOL and recovery of individuals surviving a critical illness. The rehabilitation programme for this cohort reflects similar successful programmes applied in patients with cardiac and respiratory disease [17,18] by optimizing functional recovery, particularly during the first few months after such an illness. However, there is a dearth of literature examining similar programmes involving the general critical care population [3,6,15].

## Materials and methods

This multicentre randomized controlled trial (RCT) tests the effect on physical and psychological function of an eight-week home-based rehabilitation programme for individuals who have survived a critical illness. Physical function is measured using the Physical Functioning (PF) subscale of the Short-Form-36 version 2 (SF-36) Health Survey [19] and the 6 minute walk test (6 MWT) [20]; psychological function is measured using the Depression Anxiety and Stress Scale (DASS-21) [21] and the Impact of Events Scale (IES) [22]. Other aspects of HRQOL are measured using the SF-36. The recruitment frame for the study currently includes three tertiary referral hospitals and three metropolitan hospitals in Sydney, New South Wales, and two tertiary referral hospitals and a metropolitan hospital in Brisbane, Queensland, Australia.

## Research hypothesis and aims

The primary research hypothesis is that survivors of a critical illness who participate in the physical rehabilitation programme will have better physical function, as measured by a

difference of 10 points on the SF-36 PF subscale, when compared with those who receive usual care at eight weeks after hospital discharge (short-term effect), which will persist at 26 weeks (long-term effect).

The secondary aims of the study test the effectiveness of the rehabilitation programme in terms of the following: improvement in other domains of HRQOL, as measured using the SF-36 [19]; better physical endurance, as measured by the 6 MWT [20]; improved psychological recovery, as measured using the DASS-21 [21]; less psychological distress following critical illness, as measured using the IES [22]; and reduced use of health services.

We do not propose to test *a priori* hypotheses about these secondary outcomes, because there is insufficient evidence available about the effect of the intervention on these aspects of health and well-being in this patient population and context. Rather, the study will yield information useful for formulating future hypotheses about these secondary outcomes.


## Sample size

Sample size was calculated for the SF-36 PF subscale for a two-sided hypothesis test with a type I error rate of 0.05 and a type II error rate of 0.20 (80% power). The clinically important difference and the standard deviation estimates used in our sample size calculations were based on our pilot data [23,24] and reports for similar cohorts and contexts [6,19,25,26]. At baseline we anticipate that both groups will have mean PF scores of 45. We postulate that the control group will improve by 5 points at 8 weeks, with the intervention group improving by 15 points, giving a difference of 10 points between the two study groups. Using the 10 items that comprise the PF subscale of SF-36, an improvement of 15% represents a change from 'limited a lot' to 'limited a little' on three items in the scale, for example in climbing stairs or walking particular distances. These changes reflect significant clinical improvement in physical function [27].

A sample of 100 patients per study group is required to detect this difference, assuming similar group variance (standard deviation = 25) [23,24]. We will over-enrol by 20% to account for losses to follow-up (10% study attrition [6]; 10% mortality at 6 months after hospital discharge following a critical illness [7,28]). Reasons for loss to follow up will be recorded (such as, death, withdrawal). We will minimize avoidable loss to follow up at 6 months by maintaining monthly phone contact with all participants during the study period. We will therefore enrol up to 240 participants across the recruitment sites over an 18-month recruitment period.

## Procedure

Approval to conduct the study was obtained from the Human Research Ethics Committees of the hospitals acting as recruitment sites and the universities of the investigators. The study

**Figure 1**

**Participant enrolment and retention flow diagram.** <sup>1</sup>The inclusion criteria are as follows: age  $\geq 18$  years or older; intensive care unit length of stay  $>48$  hours; receipt of mechanical ventilation for  $\geq 24$  hours; discharged home to self-care or carer; residence within the hospitals' local catchment areas to enable home visits; no neurological, spinal or skeletal dysfunction preventing participation in physical rehabilitation; not receiving palliative care; and no organised rehabilitation related to ongoing chronic disease.

protocol is registered with the Australian Clinical Trials Registry (ACTRN12605000166673) [29]. The planned flow of patients through the study reflects recommendations from the CONSORT (Consolidated Standards of Reporting Trials) statement [30], and is illustrated in Figure 1.

#### *Inclusion criteria*

To be eligible for enrolment, participants must be aged 18 years of age or older; have an ICU length of stay greater than 48 hours; have received mechanical ventilation for 24 hours or longer; be discharged home to self-care or carer (non-institutional care); reside within the hospitals' local geographical areas to enable home visits (approximately 30 km radius); have

no neurological, spinal, or skeletal dysfunction preventing participation in physical rehabilitation; not be receiving palliative care; and have no organized rehabilitation related to ongoing chronic disease management (such as, pulmonary rehabilitation, cardiac rehabilitation).

#### *Recruitment and randomization*

Eligible patients are approached immediately before or after ICU discharge, and informed voluntary consent obtained. After participant consent, the site project officer contacts an independent telephone randomization service for the participant study number and group allocation. The service uses the blocked random allocation sequences (one for each recruit-

ment site) generated using SAS software (SAS Institute Inc., Cary, NC, USA) by the study statistician (MTK), who is the only member of the study team with prospective knowledge of these sequences. Effectiveness of the randomization process will be examined by an independent Data Monitoring Committee (see Data management and analysis, below).

### *Assessment*

All participants are assessed at home within 1 week of hospital discharge by an assessor who is blinded to group assignment (Table 1). The control group receives usual care after hospital discharge. The three assessment visits for the control group are additional to 'usual care'. Although this contact is unavoidable and may have a placebo effect, any effect will reduce the apparent effectiveness of the intervention in this RCT. The treatment effect is therefore measured relative to the control group in the study, although in reality the comparator would be usual care without assessment contact.

Following the first assessor visit, participants allocated to the intervention group are visited at home by a dedicated trainer in weeks 1, 3 and 6 to provide individualized verbal and written instructions on their planned exercise programme. Exercise participants are telephoned by the trainer in weeks 2, 4, 5 and 7 to monitor their progress. Home-based follow-up assessments occur at 8 and 26 weeks after discharge for all participants (Table 1).

### *Researcher safety protocol*

A safety protocol is used to ensure assessor and trainer safety during home visits: a list of visits is held by the responsible investigator for each recruitment site with the participant's address, date, time and approximate duration of each visit; the assessor/trainer carry a mobile phone for contact; the assessor/trainer sends a text message to the investigator before entering the home and when the visit is complete; and if no contact is made, the investigator phones the assessor/trainer and then the participant's home phone (if connected).

### **Outcome measures**

The primary outcome measure is the physical functioning of study participants. We selected the SF-36 PF subscale as the corresponding measuring instrument [19] because it has demonstrated reliability, validity and responsiveness in the post-ICU population [31], and is the instrument most commonly used to assess health status in this patient cohort [3,8-10]; this will allow comparison between this study and similar cohorts.

The secondary outcome measures are as follows: exercise capacity; HRQOL; and constructs of psychological well being. Exercise capacity is measured using the 6 MWT [20]. The 6 MWT is performed twice at each assessment to account for any learning effect, with the best result recorded for analysis. During the 6 MWT, participants are monitored

continuously using a portable pulse oximeter (measuring pulse rate and oxygen saturation), with their exertion level assessed and documented during the test [20] using the Borg perceived exertion scale [32]. HRQOL is assessed using other subscales of the SF-36 [19], including effect of Physical Health on Role (Role – Physical), Bodily Pain, General Health, Vitality, Social Functioning, effect of Emotional Health on Role (Role – Emotional), and Mental Health.

We also measure constructs of psychological well being as important aspects of outcome in this patient group [3,15]. Previous studies have demonstrated that psychological distress (anxiety, depression, worry) during recovery correlate with an increased incidence of symptoms consistent with post-traumatic stress [6], although any causal relationships between critical illness, physical recovery and psychological distress are not yet clear. Psychological state is assessed using DASS-21 [21] and the IES [22]. The DASS-21 is a psychometrically robust measure of affect, using 21 items to examine the three psychological dimensions of depression, anxiety and stress, described by physical symptoms, cognitions or feelings, and rated on a 4-point scale over the preceding week. Australian normative data are available for comparison. The IES uses 15 items to measure avoidance behaviour and intrusive thoughts associated with psychological distress consistent with symptoms of post-traumatic stress disorder.

In addition, participants record all their contacts with health services during the study period, including the number and purpose of visits to their family doctor, medical specialist, community nurse, pharmacist, community health centre, or emergency department, or hospitalization since the index hospital episode or previous assessment. The battery of questionnaires takes 20–30 minutes to complete.

### **Intervention**

Participants randomized to the intervention group receive an eight-week, home-based physical rehabilitation programme that focuses on strength training and walking. All exercises use standard approaches for improving muscle strength and endurance within a rehabilitation setting [33,34]. Participants are visited at home in week 1 for exercise prescription and supervised training. Physical rehabilitation training involves graded, individualized endurance and strength training prescribed by a pulmonary rehabilitation physiotherapist. Training initially focuses on lower limb exercises and walking, with warm-up stretches. As participants progress, core stabilization and upper limb exercises are introduced. Participants are also visited by the trainer in weeks 3 and 6 for assessment of progress and reinforcement of the exercise programme, and are contacted by the trainer by telephone in nonvisit weeks (2, 4, 5, 7 and 8) to reinforce the exercise programme.

**Table 1****Participant activities and study contact**

Participant activities	Home visits after discharge (weeks) <sup>a</sup>				
	1 (baseline)	3	6	8 (short term)	26 (long term)
Outcome assessment: control and intervention groups					
Health-related quality of life (SF-36)	Visit 1	-	-	Visit 5	Visit 6
Physical endurance (6 MWT/Borg)	Visit 1	-	-	Visit 5	Visit 6
Psychological status (DASS/IES)	Visit 1	-	-	Visit 5	Visit 6
Intervention group					
Strength training	Visit 2	Visit 3	Visit 4	-	-
Endurance training	Visit 2	Visit 3	Visit 4	-	-

<sup>a</sup> Visits 1, 5 and 6 are conducted by an assessor blinded to the study group; visits 2, 3 and 4 are conducted by a 'trainer' not involved in assessment, with follow up phone calls during weeks 2, 3, 5, 7 and 8 of the programme. Borg, modified Borg Scale; DASS, Depression, Anxiety, and Stress Scale; IES, Impact of Events Scale; 6 MWT, 6 min walk test; SF-36, Short-Form-36 version 2.

**Exercise manual**

An illustrated exercise manual supports the participant's training and graded progression. The manual contains three parts: 'how hard should you exercise', which details the modified Borg Scale and provides information about participant safety; the exercise program; and how to progress the exercises. The exercise program details exercises in five components: endurance exercise (walking), lower and upper limb strengthening, core stabilization, flexibility, and stretches. A total of 16 different exercises are numbered, named, illustrated and described, to facilitate participant-trainer communication and exercise progression. This includes four stretching, three flexion, and three core stabilization exercises, which are included in the exercise prescription based on assessment of the participant's capabilities and needs by the exercise trainer.

**Strength training**

Assessment of strength uses an eight-repetition maximum (8 RM) protocol (that is to say the exercise or weight that can only be completed eight times in one set). An 8 RM protocol is less likely to cause undue strain for the participant than other assessments of strength that require heavier weights. Strength training includes upper (biceps, triceps, shoulder abductors/adductors) and lower limb (quadriceps, hamstrings, hip abductors and extensors) muscle groups. The initial strength training prescription is one set of 8 RM for each activity, progressing to three sets. Further progression is based on increasing weight (0.25–1.5 kg for arm exercises using food cans or bags of rice, or increasing the step height or weight for lower limb exercises), with levels of progression described.

**Endurance training**

Exercise prescription for endurance training is based on the results of the 6 MWT during the baseline assessment visit.

Training intensity commences at 80% of peak walking speed. Extra activities are prescribed based on a level of perceived exertion of moderate intensity (3–4, modified Borg scale). The programme uses a walk-rest-walk approach, with the duration of walking varying according to the participant's ability and condition; 12 levels of walking are described ranging from 1 to 60 minutes. Participants work toward an optimal goal of training on 5 days per week for 20–30 minutes by the end of the programme.

**Data management and analysis**

Data are entered into a purpose-built Access database at the three coordinating sites; monthly site reports on enrolment, randomization and participant follow up are submitted to one central site and monthly summaries of the information shown in Figure 1 for the whole study are given to all investigators. Analysis is by intention to treat. Analysis of covariance of the PF subscale scores for the control and intervention groups at 8 weeks will be conducted, using the baseline measure as a covariate. Multivariate regression techniques will describe the effect of the intervention on the secondary outcome measures (other SF-36 domains, 6 MWT, DASS subscales and IES subscales), and correlations among all outcome measures will be estimated. Baseline characteristics of patients lost to follow up at 8 weeks and 26 weeks will be compared with patients who completed follow up to assess patterns of loss to follow up and provide insights into generalizability of the results. A Data Monitoring Committee independent of the research team will be convened to review study progress and interim and final analyses [35].

**Pilot study**

The intervention and all study procedures were tested in a small dual-site pilot RCT before commencement of the fully powered study described here. The pilot study demonstrated

the feasibility of the training programme and study procedures (recruitment, retention, outcome measures), and enabled refinement of the training manual, researcher safety procedures, protocol implementation across multiple sites, and data collection and management strategies.

## Results and discussion

The burden for survivors of a critical illness has been well documented in many observational studies, in which the recovery trajectory is often prolonged and suboptimal. However, intervention studies with this clinical cohort are less common. To our knowledge, the proposed research is the first study internationally to use a home-based rehabilitation programme in this patient group. An individualized, home-based programme negates the need to attend an outpatient clinic located in a hospital on a regular basis. This is particularly important for individuals who reside in regional or rural areas but were treated in a metropolitan ICU, as well as those who choose not to or are unable to participate in hospital-based programmes for other reasons such as lack of mobility or transport. The provision of such a programme through local community health services would allow survivors of a critical illness to engage in the programme regardless of place of residence and other mobility and access constraints.

The significance of this research can be considered from a number of aspects. The first aspect pertains to the patient and their family. The effects of a critical illness can be devastating on all aspects of HRQOL, including physical, psychological, social, fiscal and spiritual. This intervention aims to improve the recovery for patients to their optimal level of functioning following a critical illness using a structured programme of physical exercise. The second aspect concerns health professionals. Following the patient through their recovery process will help to inform clinicians regarding the appropriateness and long-term impact of ICU management. Development of outreach services to ensure any benefits gained are not lost by insufficient support after intensive care is an important component of clinical management within an extended episode of care. Finally, the research findings will be of value to health service managers because they may facilitate utilization of health care resources appropriately and efficiently to improve patient outcomes for survivors of a critical illness.

If the study findings are positive and conclusive, this approach could be adopted by any community or primary health service in both metropolitan and rural settings, given appropriate training and support. The exercise manual uses standard exercises and training patterns. Facilitation and monitoring of individuals engaged in the programme could be undertaken by a variety of health professionals within primary care or community health services (for example, general practitioners, community nurses, practice nurses, physiotherapists) after some initial training and with the aid of resources such as the training manual.

The innovative aspects of the project are in their application to this important but often heterogeneous group of patients and in the project's home-based focus, which frees patients from the need to attend hospital outpatient clinics. This research will also inform clinicians and health service providers about one approach within a range of outreach services that best serve the needs of patients following critical illness and lead to better long-term patient outcomes with appropriate utilization of scarce resources.

## Conclusion

This study uses a multicentre RCT design to provide the best evidence to examine the effectiveness of a novel application of physical rehabilitation practices to an important clinical cohort. The project's significance relates to two key features. First, the study addresses outcomes that are meaningful for patients and society, namely functional ability and well being. Second, it is targeted at a health problem that is likely to increase as the population ages, contributing to an area in which there are minimal rigorous intervention studies. If the programme is effective, it will provide a model that can be easily adapted and adopted by existing primary care or community services.

### Key messages

- Many observational studies and associated systematic reviews have demonstrated delayed physical and psychological recovery for survivors of a critical illness, often beyond 6 months after discharge.
- There have been limited interventional studies aimed at improving the recovery process for this patient cohort.
- This novel RCT with blinded assessment tests the effects of an eight-week, home-based, individually tailored physical rehabilitation programme on physical and psychological recovery for survivors of a critical illness after discharge from hospital.
- Participants in the intervention group receive graded, individualized endurance and strength training, focusing initially on lower limb exercises and walking, with warm up stretches, and progresses to the addition of core stabilization and upper limb exercises.
- Participant outcomes, including physical endurance, HRQOL, and psychological well being (depression, anxiety, stress, avoidance, intrusive thoughts) will be assessed using standard instruments at weeks 1, 8 and 26 after hospital discharge.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

All authors contributed to the study design and methods, and the development of the grant application that was the source for this paper. JA specifically contributed to development of

the exercise manual. MK, SM and DE specifically contributed to the statistical methods and power calculations. DE drafted the manuscript and all other authors critically revised it for important intellectual content. All authors approved the final version of the manuscript for publication.

## Acknowledgements

This research is supported by the National Health and Medical Research Council (NHMRC, Australia; Project Grant 352407). Preliminary work was supported by a seed grant from Effective Healthcare Australia, and a Sesqui Research and Development Grant from The University of Sydney. The authors thank Harriet Adamson and Anne-Marie Boxall for their initial involvement in the preliminary work for this project. Thanks also to the current project staff, Penny Taylor, Rachel Foley, Jacelle Lang, Naomi Hammond and Linda Parker, and all the clinical staff involved in patient screening and participant recruitment activities.

## References

- Black NA, Jenkinson C, Hayes JA, Young D, Vella K, Rowan KM, Daly K, Ridley S: **Review of outcome measures used in adult critical care.** *Crit Care Med* 2001, **29**:2119-2124.
- Needham DM, Dowdy DW, Mendez-Tellez PA, Herridge MS, Pronovost P: **Studying outcomes of intensive care unit survivors: measuring exposures and outcomes.** *Intensive Care Med* 2005, **31**:1153-1160.
- Angus DC, Carlet J: **Surviving intensive care: a report from the 2002 Brussels Roundtable.** *Intensive Care Med* 2003, **29**:368-377.
- Ball C, Kirkby M, Williams S: **Effect of the critical care outreach team on patient survival to discharge from hospital and readmission to critical care: non-randomised population based study.** *BMJ* 2003, **327**:1014.
- Endacott R, Chaboyer W: **The nursing role in ICU outreach: an international exploratory study.** *Nurs Crit Care* 2006, **11**:94-102.
- Jones C, Skirrow P, Griffiths RD, Humphris GH, Ingleby S, Eddleston J, Waldmann C, Gager M: **Rehabilitation after critical illness: a randomized, controlled trial.** *Crit Care Med* 2003, **31**:2456-2461.
- Higlett T, Bishop N, Hart GK, Hicks P: *Review of Intensive Care Resources & Activity 2002/2003* Melbourne: Australian and New Zealand Intensive Care Society; 2005.
- Adamson H, Elliott D: **Quality of life after a critical illness: a review of general ICU studies 1998-2005.** *Aust Crit Care* 2005, **18**:50-60.
- Elliott D: **Measuring the health outcomes of general ICU patients: a systematic review of methods and findings.** *Aust Crit Care* 1999, **12**:132-140.
- Dowdy DW, Eid MP, Sedrakyan A, Mendez-Tellez PA, Pronovost PJ, Herridge MS, Needham DM: **Quality of life in adult survivors of critical illness: a systematic review of the literature.** *Intensive Care Med* 2005, **31**:611-620.
- Herridge MS: **Long-term outcomes after critical illness.** *Curr Opin Crit Care* 2002, **8**:331-336.
- Cuthbertson BH, Hull A, Strachan M, Scott J: **Post-traumatic stress disorder after critical illness requiring general intensive care.** *Intensive Care Med* 2004, **30**:450-455.
- Jackson JC, Gordon MW, Ely EW, Burger C, Hopkins RO: **Research issues in the evaluation of cognitive impairment in intensive care unit survivors.** *Intensive Care Med* 2004, **30**:2009-2016.
- Heyland DK, Guyatt G, Cook DJ, Meade M, Juniper E, Cronin L, Gafni A: **Frequency and methodologic rigor of quality-of-life assessments in the critical care literature.** *Crit Care Med* 1998, **26**:591-598.
- Rubenfeld GD, Curtis JR: **Health status after critical illness: Beyond descriptive studies.** *Intensive Care Med* 2003, **29**:1626-1628.
- Martin UJ, Hincapie L, Numchuk M, Gaughan J, Criner GJ: **Impact of whole-body rehabilitation in patients receiving chronic mechanical ventilation.** *Crit Care Med* 2005, **33**:2259-2265.
- Oh EG: **The effects of home-based pulmonary rehabilitation in patients with chronic lung disease.** *Int J Nurs Stud* 2003, **40**:873-879.
- Boxall A, Barclay L, Sayers A, Caplan G: **A randomised controlled trial of home-based pulmonary rehabilitation for COPD patients aged over 60 years.** *Thorax* 2003, **58**:Siii76.
- Ware JE, Snow KK, Kosinski M: *SF-36 Version 2 Health Survey: Manual and Interpretation Guide* Lincoln: Quality Metric Incorporated; 2000.
- American Thoracic Society: **Guidelines for the six-minute walk test.** *Am J Respir Crit Care Med* 2002, **166**:111-117.
- Lovibond P, Lovibond S: **The structure of negative emotional states: comparison of the Depression Anxiety Stress Scale (DASS) with the Beck Depression and Anxiety Inventories.** *Behav Res Ther* 1995, **33**:335-343.
- Horowitz M, Wilner N, Alvarez W: **Impact of events scale: a measure of subjective stress.** *Psychosom Med* 1979, **41**:209-218.
- Kelly M, McKinley S: **Patients' physical problems in early recovery after intensive care.** *Aust Crit Care* 2003, **16**:29-30.
- Boyle M, Murgo M, Adamson H, Gill J, Elliott D, Crawford M: **The effect of chronic pain on health related quality of life amongst intensive care survivors.** *Aust Crit Care* 2004, **17**:104-113.
- Schelling G, Richter M, Roozendaal B, Rothenhausler HB, Krause-neck T, Stoll C, Nollert G, Schmidt M, Kapfhammer HP: **Exposure to high stress in the intensive care unit may have negative effects on health-related quality-of-life outcomes after cardiac surgery.** *Crit Care Med* 2003, **31**:1971-1980.
- Schelling G, Stoll C, Haller M, Briegel J, Manert W, Hummel T, Lenthart A, Heyduck M, Polasek J, Meier M, et al.: **Health-related quality of life and posttraumatic stress disorder in survivors of the acute respiratory distress syndrome.** *Crit Care Med* 1998, **26**:651-659.
- Needham DM, Dennison CR, Dowdy DW, Mendez-Tellez PA, Ciesla N, Desai SV, Sevransky J, Shanholtz C, Scharfstein D, Herridge MS, et al.: **Study protocol: The improving care of acute lung injury patients (ICAP) study.** *Crit Care* 2005, **10**:R9.
- Chelluri L, Im KA, Belle SH, Schulz R, Rotondi AJ, Donahoe MP, Sirio CA, Mendelsohn AB, Pinsky MR: **Long-term mortality and quality of life after prolonged mechanical ventilation.** *Crit Care Med* 2004, **32**:61-69.
- Australian Clinical Trials Registry** [<http://www.actr.org.au>]
- Moher D, Schulz KF, Altman DG, for the CONSORT Group: **The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomised trials.** *Lancet* 2001, **357**:1191-1194.
- Chrispin PS, Scotton H, Rogers J, Lloyd D, Ridley SA: **Short form 36 in the intensive care unit: assessment of acceptability, reliability and validity of the questionnaire.** *Anaesthesia* 1997, **52**:15-23.
- Borg GA: **Psychophysical bases of perceived exertion.** *Med Sci Sports Exerc* 1982, **14**:377-381.
- Barker-Whittle Z, Alison J, Bye P: **Arm and leg exercise capacity in subjects with severe CAL compared with healthy control subjects.** *Respirology* 2002, **7s**:A33.
- Alison J, Spencer L: **Effects of pulmonary rehabilitation on 12 minute walk distance in lung volume reduction surgery patients.** *Respirology* 2000, **5S**:A44.
- DAMOCLES Study Group: **A proposed charter for clinical trial data monitoring committees: helping them do their job well.** *Lancet* 2005, **365**:711-722.